


# XYZ States at BESIII

Shan Gu 顾珊

Beihang University && IHEP, Beijing



( for the BESIII Collaboration )


**RENCONTRES DU VIETNAM**  
**25th Anniversary**

Quy Nhon, August 8, 2018

# Outline

- The BESIII experiment
- Charmoniumlike states
  - The Y states —  $J^{PC}=1^{--}$   
 $Y \rightarrow \pi^+ \pi^- J/\psi$ ,  $Y \rightarrow \pi^+ \pi^- \psi(3686)$ ,  $Y \rightarrow \pi^+ \pi^- h_c$
  - The  $Z_c$  states —  $I=1$  & decays into  $c\bar{c}$   
 $J^P$  for  $Z_c(3900)$ ,  $Z_c \rightarrow \rho \eta_c$
  - The  $X(3872)$  —  $J^{pc}=1^{++}$   
 $e^+ e^- \rightarrow \gamma X(3872)$ ,  $X(3872) \rightarrow \pi^+ \pi^- J/\psi$
- Summary

# Beijing Electron Positron Collider II (BEPCII)




# Beijing Spectrometer (BESIII)


Nucl. Instr. Meth. A 614, 345 (2010)

## □ Inner to Outside:

- Main Drift chamber(MDC)
- Time of flight System(TOF)
- Electromagnetic Calorimeter(EMC)
- Solenoid super-conducting magnet(SSM)
- Muon chamber(MUC)





# BESIII data sets for XYZ study



# The Y states

# Precise Measurement of $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ Cross Section



PRL118, 092001 (2017)



- Simultaneous fit to the cross section from XYZ data(left) and R-scan data (right)
- Fit I =  $|BW1 + BW2 * e^{i\phi^2} + BW3 * e^{i\phi^3}|^2$  or Fit II =  $|\exp + BW2 * e^{i\phi^2} + BW3 * e^{i\phi^3}|^2$
- Resonance I:  $M = (4222.0 \pm 3.1 \pm 1.4) \text{ MeV}$ ,  $\Gamma = (44.1 \pm 4.3 \pm 2.0) \text{ MeV}$ ,  
**Lower and narrower than previous Y(4260) PDG value**
- Resonance II:  $M = (4320.0 \pm 10.4 \pm 7) \text{ MeV}$ ,  $\Gamma = (101.4 \pm 25 \pm 10) \text{ MeV}$ ,  
**a little bit lower than Y(4360) PDG value**
- The significance of the second resonance is  $7.6\sigma$
- The first observation of  $Y(4360) \rightarrow \pi^+\pi^-J/\psi$ ?

$$e^+ e^- \rightarrow \pi^+ \pi^- \psi(3686)$$

## ➤ Reconstructed modes


PRD 96, 032004 (2017)

- **Mode I:**  $\psi(3686) \rightarrow \pi^+ \pi^- \text{J}/\psi$ ,  $\text{J}/\psi \rightarrow l^+ l^- (l = e/\mu)$ .
  - **Mode II:**  $\psi(3686) \rightarrow neutrals + \text{J}/\psi$ ,  
 $neutrals = (\pi^0 \pi^0, \pi^0, \eta \text{ and } \gamma\gamma)$ ,  $\text{J}/\psi \rightarrow l^+ l^- (l = e/\mu)$ .



Ecm=4.416 GeV

# $e^+e^- \rightarrow \pi^+ \pi^- \psi(3686)$




PRD 96, 032004 (2017)

Parameters	Solution I	Solution II
$M(Y4220)$ (MeV/c <sup>2</sup> )	$4209.5 \pm 7.4$	
$\Gamma(Y(4220))$ (MeV)	$80.1 \pm 24.6$	
$\mathcal{B}\Gamma^{e^+e^-}(Y(4220))$ (eV)	$0.8 \pm 0.7$	$0.4 \pm 0.3$
$M(Y4390)$ (MeV/c <sup>2</sup> )	$4383.8 \pm 4.2$	
$\Gamma(Y(4390))$ (MeV)	$84.2 \pm 12.5$	
$\mathcal{B}\Gamma^{e^+e^-}(Y(4390))$ (eV)	$3.6 \pm 1.5$	$2.7 \pm 1.0$
$\phi_1$ (rad)	$3.3 \pm 1.0$	$2.8 \pm 0.4$
$\phi_2$ (rad)	$0.8 \pm 0.9$	$4.7 \pm 0.1$

- Fix parameters of the  $Y(4660)$  to Belle results.
- The  $Y(4220)$  is necessary(significance =  $5.8\sigma$  )

$$e^+ e^- \rightarrow \pi^+ \pi^- h_c$$

PRL118, 092002 (2017)



The cross sections are of the same order of magnitude as those of the  $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi(\psi(3686))$ , but follow a different line shape.





- Fitted with coherent sum of two Breit-Wigner like structure

$$M_1 = 4218.4^{+5.5}_{-4.5} \pm 0.9 \text{ MeV}/c^2, \Gamma_1 = 66.0^{+12.3}_{-8.3} \pm 0.4 \text{ MeV} \rightarrow Y(4220)$$

$$M_2 = 4391.5^{+6.3}_{-6.8} \pm 1.0 \text{ MeV}/c^2, \Gamma_2 = 139.5^{+16.2}_{-20.6} \pm 0.6 \text{ MeV} \rightarrow Y(4390)$$

# The $Z_c$ states


# Determination of $J^p$ of Zc(3900)




PRL 119, 072001 (2017)

- Based on a PWA of the  $e^+e^- \rightarrow \pi^+\pi^- J/\psi$  at  $\sqrt{S} = 4.23, 4.26$  GeV.
- Assume Zc to have  $J^p=1^+$ , parameterized with Flatte formula.
- Pole Mass =  $(3881.2 \pm 4.2 \pm 52.7)$  MeV,  
Pole width =  $(51.8 \pm 4.6 \pm 36.0)$  MeV

# Determination of $J^P$ of Zc(3900)




Polar angle distribution of  $Z_c^\pm$  in the process  $e^+e^- \rightarrow Zc^\pm\pi^- + \text{c.c.}$




Helicity angle distribution of  $J/\psi$  in the  $Zc^\pm \rightarrow \pi^\pm J/\psi$ .

The significance of the  $J^P=1^+$  hypothesis over the alternative  $J^P$  possibilities to be larger than  $7\sigma$ .

# Search for $Z_c \rightarrow \rho\eta_c$



$$R_z = \frac{B(Z_c \rightarrow \rho\eta_c)}{B(Z_c \rightarrow \pi J/\psi)}$$



$$R_{z'} = \frac{B(Z'_c \rightarrow \rho\eta_c)}{B(Z'_c \rightarrow \pi h_c)}$$

*A. Esposito, A.L. Guerrieri, A. Pilloni, Phys. Lett. B 746, 194 (2015)*

- The ratios of  $Z_c^{(\prime)} \rightarrow \rho\eta_c$  to  $Z_c^{(\prime)} \rightarrow \pi J/\psi(\pi h_c)$  may distinguish the tetra-quark and molecule models.
- The green band and yellow band show the  $1\sigma$  and  $2\sigma$  confidence range of the corresponding theoretical model.


# Evidence for $Z_c \rightarrow \rho\eta_c$

preliminary

- $e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta_c$
- $\eta_c \rightarrow 9$  hadronic decays

Decay mode	BR
$\eta_c \rightarrow p\bar{p}$	$\sim 0.13\%$
$\eta_c \rightarrow 2(K^+K^-)$	$\sim 0.15\%$
$\eta_c \rightarrow \pi^+\pi^-K^+K^-$	$\sim 1.50\%$
$\eta_c \rightarrow K^+K^-\pi^0$	$\sim 1.20\%$
$\eta_c \rightarrow p\bar{p}\pi^0$	$\sim 0.18\%$
$\eta_c \rightarrow K_SK\pi$	$\sim 1.80\%$
$\eta_c \rightarrow \pi^+\pi^-\eta$	$\sim 1.60\%$
$\eta_c \rightarrow K^+K^-\eta$	$\sim 0.57\%$
$\eta_c \rightarrow \pi^+\pi^-\pi^0\pi^0$	$\sim 2.40\%$

- Strong evidence for  $e^+e^- \rightarrow \pi Z_c, Z_c \rightarrow \rho\eta_c$  at  $\sqrt{s} = 4.23\text{GeV}$   
Significance:  $4.3\sigma$ .
- $e^+e^- \rightarrow \pi Z'_c, Z'_c \rightarrow \rho\eta_c$  not seen.  
Significance:  $1\sigma$ .




$e^+e^- \rightarrow \pi Z_c, Z_c \rightarrow \rho\eta_c$  at  $\sqrt{s} = 4.23\text{GeV}$

# Evidence for $Z_c \rightarrow \rho\eta_c$


preliminary

	$\sqrt{s} = 4.23 \text{ GeV}$	$\sqrt{s} = 4.26 \text{ GeV}$	$\sqrt{s} = 4.36 \text{ GeV}$	Tetra-quarks-I	Tetra-quarks-II	Molecule
$R_{Z_c(3900)}$	$2.1 \pm 0.8$	$< 6.4$	...	$230^{+330}_{-140}$	$0.27^{+0.40}_{-0.17}$	$0.046^{+0.025}_{-0.017}$
$R_{Z_c(4020)}$	$< 1.9$	$< 1.2$	$< 1.0$		$6.6^{+56.8}_{-5.8}$	$0.010^{+0.006}_{-0.004}$



$$R_z = \frac{B(Z_c \rightarrow \rho\eta_c)}{B(Z_c \rightarrow \pi J/\psi)}$$

A. Esposito et al, Phys. Lett. B 746, 194 (2015)



$$R_{z'} = \frac{B(Z'_c \rightarrow \rho\eta_c)}{B(Z'_c \rightarrow \pi h_c)}$$

Also calculations predict  
very different values:

$$R_z = 10^{-3} \sim 10^2 !$$

arXiv:1806.05651

arXiv:1512.01938

PRD 91, 034032 (2015)


PRD 90, 054006 (2014)

EPJC 73, 2561 (2013)


# The X state

# $e^+e^- \rightarrow \gamma X(3872), X(3872) \rightarrow \pi^+ \pi^- J/\psi$

PRL 112, 092001 (2014)



- The  $X(3872)$  signal is clearly observed: **significance  $6.3\sigma$ .**  
 $M(X(3872)) = 3871.9 \pm 0.7 \pm 0.2 \text{ MeV}$  [PDG:  $3871.68 \pm 0.17 \text{ MeV}$ ]
- Existence of radiative transition process  
 $Y(4260) \rightarrow \gamma X(3872)$ .
- Assuming that measured transition is from  $Y(4260)$  and  $\mathcal{B}[X(3872) \rightarrow \pi^+ \pi^- J/\psi] = 5\%$



$$\frac{\mathcal{B}[Y(4260) \rightarrow \gamma X(3872)]}{\mathcal{B}(Y(4260) \rightarrow \pi^+ \pi^- J/\psi)} = 0.1$$

# Summary

- BESIII collaboration has performed a detailed study of the XYZ at 3.8-4.6 GeV
  - **Measurement of Born cross-section for different channels reveal complex structures and new Y states**
  - **$J^P=1^+$  for  $Z_c(3900)$ , evidence for  $Z_c \rightarrow \rho\eta_c$**
  - **Observation of  $Y(4260) \rightarrow \gamma X(3872)$ .**
- BESIII will take more data and continue the study.

**Thanks!!!**