


# Determination of the Number of Neutrinos at LEP (30 years after)

Monica Pepe Altarelli (CERN)

XVth Rencontres du Vietnam, “3 neutrinos and Beyond”  
4-10 August 2019, ICISE, Qui Nhon



### Announcement from the Director-General

I am pleased to announce to you that the LEP machine is operating at 45.5 GeV and that Z<sup>0</sup> particles have been observed.

This is a great achievement of which we can all be justly proud, especially the many of you who have made it all possible.


Carlo Rubbia  
Director-General  
14 August 1989

### Communication du Directeur général


Je suis heureux de vous annoncer que la machine LEP fonctionne à 45,5 GeV et que des particules Z<sup>0</sup> y ont été observées.

C'est un grand succès dont nous pouvons tous être justement fiers, plus particulièrement les nombreuses personnes parmi vous qui l'ont rendu possible.


Carlo Rubbia  
Directeur général  
le 14 août 1989




DELPHI



ALEPH



L3



OPAL

### Z<sup>0</sup> marks the spot

Late on the night of Sunday 13 August, just one month after first beam circulated and a mere 16 minutes after the start of the pilot run, LEP's first Z<sup>0</sup> was recorded. By midnight a total of three had been observed, and on Monday there followed 13 more - a remarkable total of 16 between the four detectors ALEPH, DELPHI, OPAL and L3 in the first 24 hours of operation.

### Le règne du Z<sup>0</sup>

Tard dans la nuit du dimanche 13 août, un r exactement après les premières révolutions faisceaux dans l'anneau et 16 minutes seules après le début de la période d'essai, le premier Z<sup>0</sup> LEP a été enregistré. A minuit leur nombre s'éleva à trois et lundi 13 autres ont suivi, soit un total remarquable de 16 Z<sup>0</sup> pour l'ensemble des quatre détecteurs ALEPH, DELPHI, OPAL et L3 au cours des 24 premières heures d'exploitation.

# Let us turn our clock back



# What was known at the time?

- SM of electroweak interactions well established with the W and Z discovery in 1983 and the measurement of their properties
- SM however does not predict the number of fermion generations or their masses

- Quarks and leptons organised into three families →

	I	II	III
Quarks	$2/3$ Left Right 2.4 MeV <b>u</b> up	$2/3$ Left Right 1.27 GeV <b>c</b> charm	$2/3$ Left Right 173.2 GeV <b>t</b> top
	$-1/3$ Left Right 4.8 MeV <b>d</b> down	$-1/3$ Left Right 104 MeV <b>s</b> strange	$-1/3$ Left Right 4.2 GeV <b>b</b> bottom
	$0$ Left Right <b><math>\nu_e</math></b> electron neutrino	$0$ Left Right <b><math>\nu_\mu</math></b> muon neutrino	$0$ Left Right <b><math>\nu_\tau</math></b> tau neutrino
Leptons	$-1$ Left Right 0.511 MeV <b>e</b> electron	$-1$ Left Right 105.7 MeV <b><math>\mu</math></b> muon	$-1$ Left Right 1.777 GeV <b><math>\tau</math></b> tau

- Basic question was:

**Are there more families than the three observed so far?**

- Given the regularity of the pattern, counting the number of neutrino types  $N_\nu$  may also mean counting the number of fundamental fermion generations

# What was known at the time? (II)


## Info on $N_\nu$ from cosmology, astrophysics & particle physics

- **Cosmology**, from primordial nucleosynthesis:  
neutrinos enter through the reaction  $n + \nu_e \leftrightarrow p + e^-$  ;  
formation of light elements and their relative abundances (He/H) is  
a function of  $N_\nu$
- **Astrophysics**, based on observation of  $\bar{\nu}$  emitted by SN 1987A,  
relying on theory and based on assumption that total gravitational  
energy release shared equally by all neutrino species
- **Particle physics**: indications from direct search from the process  
 $e^+e^- \rightarrow \nu\bar{\nu}\gamma$  and from Z/W properties at CERN and FERMILAB  
 $p\bar{p}$  experiments, e.g.  $\sigma(W \rightarrow l\nu)/\sigma(Z \rightarrow l\bar{l})$

# What was known at the time? (III)

Denegri, Sadoulet & Spiro  
[Rev. Mod. Phys, 1989]

- Remarkable agreement between values derived from the analysis of such widely different phenomena
- Putting everything together, Denegri, Sadoulet & Spiro obtained  $N_\nu = 2.0^{+0.6}_{-0.4}$





- “Results perfectly compatible with the *a priori* knowledge that at least three neutrino families should exist ... Although the consistency is significantly worse, four families still provide a reasonable fit”



# SLAC Linear Collider

- SLC was the prototype of a new accelerator concept, the linear collider
- Scheduled to take first data in Jan.'87, but new and difficult technology → First reasonable Lumi only in March '89 (few  $10^{27} \text{cm}^{-2} \text{s}^{-1}$ )
- First results based on 106 Z events collected at 6 different energies around the Z peak



$$M_Z = 91.11 \pm 0.23 \text{ GeV}/c^2$$

$$\Gamma_Z = 1.61^{+0.60}_{-0.43} \text{ GeV}$$

$$\Gamma_{inv} = 0.62 \pm 0.23 \text{ GeV} \rightarrow N_\nu = 3.8 \pm 1.4$$

[PRL 63, 724 (1989)]

# LEP

- LEP start-up advertised for 14 July 1989
  - July 14, First turn
  - August 13, First Collisions
  - August 13-18, Physics pilot run
  - August 21-Sept. 11, Machine studies
  - Sept. 20-Nov. 5, Physics




- The Economist August 19, 1989

“The results from California are impressive, especially as they come from a new and unique type of machine. They may provide a sure answer to the generation problem before LEP does. This explains the haste with which the finishing touches have been applied to LEP. The 27km-long device, six years in the making was transformed from inert hardware to working machine in just four weeks--- a prodigious feat, unthinkable anywhere but at CERN.....

.....Even so, it was still not as quick as Dr. Carlo Rubbia, CERN’s domineering director-general might have liked”.


**13 November 1989**  
**LEP Inauguration**



# First results from LEP

13 October 1989

- After only three weeks of data-taking, CERN seminar in which the four experiments presented their results based on  $\sim 3000$  Z each (J.Lefrancois, U.Amaldi, S.Ting, A.Wagner) & MARKII update
- Results written on blackboard by John Thresher, CERN research director with responsibility for the new LEP experimental programme



# First results from LEP

13 October 1989

- After only three weeks of data-taking, CERN seminar in which the four experiments presented their results based on  $\sim 3000$  Z each (J.Lefrancois, U.Amaldi, S.Ting, A.Wagner) & MARKII update
- Results written on blackboard by John Thresher, CERN research director with responsibility for the new LEP experimental programme

[A.Blondel arXiv:1812.11362v2]

Experiment	Hadronic Zs	Z mass (GeV)			$N_\nu$		
MARKII	450	91.14	$\pm$	0.12	2.8	$\pm$	0.60
L3	2538	91.13	$\pm$	0.06	3.42	$\pm$	0.48
ALEPH	3112	91.17	$\pm$	0.05	3.27	$\pm$	0.30
OPAL	4350	91.01	$\pm$	0.05	3.12	$\pm$	0.42
DELPHI	1066	91.06	$\pm$	0.05	2.4	$\pm$	0.64
Average		91.10	$\pm$	0.05	3.12	$\pm$	0.19

# First results from LEP

13 October 1989

- After only three weeks of data-taking, CERN seminar in which the four experiments presented their results based on  $\sim 3000$  Z each (J.Lefrancois, U.Amaldi, S.Ting, A.Wagner) & MARKII update
- Results written on blackboard by John Thresher, CERN research director with responsibility for the new LEP experimental programme

[A.Blondel arXiv:1812.11362v2]





Experiment	Hadronic Zs	Z mass (GeV)			$N_\nu$		
MARKII	450	91.14	$\pm$	0.12	2.8	$\pm$	0.60
L3	2538	91.13	$\pm$	0.06	3.42	$\pm$	0.48
ALEPH	3112	91.17	$\pm$	0.05	3.27	$\pm$	0.30
OPAL	4350	91.01	$\pm$	0.05	3.12	$\pm$	0.42
DELPHI	1066	91.06	$\pm$	0.05	2.4	$\pm$	0.64
Average		91.10	$\pm$	0.05	3.12	$\pm$	0.19

The number of light neutrinos was three!

# The four LEP experiments

- **ALEPH**: main emphasis on momentum measurement via accurate tracking in high magnetic field (1.5 Tesla); high granularity ECAL
  - Vertex detector (Silicon strips), Inner Tracking Chamber, Time Projection Chamber (main tracking detector) →  $\Delta p/p \simeq 2.7\%$  for 45 GeV muons
- **DELPHI**: pioneering new techniques
  - PID via RICH detector (with liquid and gas radiators), Heavy Projection Chamber used as electromagnetic calorimeter (excellent spatial resolution but complex to operate), Silicon Vertex detector
- **L3**: general tracking minimised in favour of precise outer tracking for muons only in very large solenoidal magnet ( $\Delta p/p \simeq 2.5\%$  for 45 GeV muons); Bismuth Germanium Oxide (BGO) electromagnetic calorimeter ( $\Delta E/E \simeq 1.4\%$  for 45 GeV electrons)
  - About twice as expensive as the other detectors!
- **OPAL**: more conservative design
  - Excellent tracking achieved by means of a “jet-type” drift chamber; Silicon Vertex detector installed in 1992

# The four LEP experiments



# Z line shape

- Cornerstone of LEP physics programme, studied by measuring the visible cross-section at several centre-of-mass energies near the Z mass

- For  $e^+e^- \rightarrow Z \rightarrow f\bar{f}$ , 
$$\sigma_f(s) = \sigma_f^0 \frac{s\Gamma_Z^2}{(s - M_Z^2)^2 + s^2 \frac{\Gamma_Z^2}{M_Z^2}} (1 + \delta_{\text{rad}}(s))$$

with the peak cross-section 
$$\sigma_f^0(s = M_Z^2) = 12\pi \frac{\Gamma_e \Gamma_f}{M_Z^2 \Gamma_Z^2}$$


- From measured cross-sections, EW parameters extracted after correcting for QED effects (ISR), which are large ( $\sim 30\%$  at the peak) but precisely known ( $5 \times 10^{-4}$ )
- Dependence on  $N_\nu$  through  $\Gamma_Z = 3\Gamma_\ell + \Gamma_{\text{had}} + N_\nu \Gamma_\nu$
- In the SM,  $\Gamma_{\text{had}} \sim 70\%$ ,  $3\Gamma_\ell \sim 10\%$  and  $\Gamma_{\text{inv}} = N_\nu \Gamma_\nu \sim 20\%$

# $\sigma(e^+e^- \rightarrow \text{hadrons})$ as a function of $\sqrt{s}$

- Drawn in 1987 before the start of LEP
- One additional  $\nu$  species would increase  $\Gamma_z$  by 6.6% and decrease the peak cross-section  $\sigma_{\text{had}}^0$  by 13%

$$\frac{d\sigma_{\text{had}}^0/dN_\nu}{\sigma_{\text{had}}^0} = -13\%$$

[A.Blondel, arXiv:1812.11362]



# The method

- Primary quantities measured :  $\sigma(\sqrt{s})$  for hadrons,  $e^+e^-$ ,  $\mu^+\mu^-$ ,  $\tau^+\tau^-$  (but first  $N_\nu$  measurement only based on hadrons)
- From  $\sigma(\sqrt{s})$  to a final state  $f\bar{f}$  one can extract peak position, width and overall normalisation, best obtained from the peak cross-section  $\sigma_f^0$
- Fit for  $M_Z$ ,  $\Gamma_Z$ ,  $\sigma_{\text{had}}^0$  and  $R_{e,\mu,\tau} \equiv \sigma_{\text{had}}^0/\sigma_{e,\mu,\tau}^0 = \Gamma_{\text{had}}/\Gamma_{e,\mu,\tau}$  (choice that minimizes experimental correlations)
- Parameter set reduces to four:  $M_Z, \Gamma_Z, \sigma_{\text{had}}^0, R_\ell$  assuming lepton universality

$$N_\nu = \frac{\Gamma_\ell}{\Gamma_\nu} \cdot \left( \sqrt{\frac{12\pi R_\ell}{M_Z^2 \sigma_{\text{had}}^0}} - R_\ell - 3 \right)$$


- Dominant sensitivity in  $N_\nu$  determination through hadronic peak cross-section  
[G.Feldman, '87]

# Principles of the analysis


- All visible Z decays detected and classified according to the four categories:  
hadrons,  $e^+e^-$ ,  $\mu^+\mu^-$ ,  $\tau^+\tau^-$

- High and well-known efficiency, e.g.  
 $\epsilon_{\text{had}} > 99\%$


$$e^+e^- \rightarrow q\bar{q}$$




$$e^+e^- \rightarrow e^+e^-$$



$$e^+e^- \rightarrow \mu^+\mu^-$$




$$e^+e^- \rightarrow \tau^+\tau^-$$



# Luminosity measurement

- Uncertainty on Luminosity has direct impact on  $N_\nu$  :  $\Delta N_\nu \sim 8 \frac{\Delta \mathcal{L}}{\mathcal{L}}$
- Luminosity determined by measuring at the same time another process with known cross-section, low-angle Bhabha scattering :  $e^+e^- \rightarrow e^+e^-$  (dominated by  $t$ -channel  $\gamma$ -exchange) through dedicated detectors for the scattered electrons



- Method: compare measured rate of Bhabha scattering with cross-section predicted by theory

$$\mathcal{L} = \frac{N_{\text{Bhabha}}}{\sigma_{\text{ref}}}$$

# events in data passing selection cuts

Calculated from MC events using same cuts as for data

# Luminosity measurement (II)

- Experimental challenge was to define the geometrical acceptance with high accuracy, especially at the lowest  $\theta$  bound:


$$\frac{d\sigma}{d\theta} \sim \frac{1}{\theta^4} \rightarrow \sigma^{\text{acc}} \sim \frac{1}{\theta_{\text{min}}^2} - \frac{1}{\theta_{\text{max}}^2} \rightarrow$$
$$\frac{\delta\sigma^{\text{acc}}}{\sigma^{\text{acc}}} \simeq \frac{2\delta\theta_{\text{min}}}{\theta_{\text{min}}} = 2 \left( \frac{\delta R_{\text{min}}}{R_{\text{min}}} \oplus \frac{\delta z}{z} \right) \quad \theta \simeq R/z$$

- ALEPH: Sensitivity to possible displacements with respect to beam position reduced with asymmetrical event selection: tight fiducial cut on one side (e.g.  $e^+$ ) and loose on the other ( $e^-$ ) with fiducial role alternating from one event to the next (plus other clever tricks  $\rightarrow$  experimental systematics  $\sim 1\%$  already in first paper)
- Experimental precision decreased to well below  $10^{-3}$  at the end of LEP after progressive replacement with more precise calorimeters, e.g. silicon-tungsten
- After a lot of hard work final precision on theory estimate of cross-section within acceptance reached  $6 \times 10^{-4}$  (from Monte Carlo program for small-angle Bhabha scattering BHLUMI) matching experimental uncertainty



# Some archeology

# Some days before LEP startup



# First ALEPH analysis approval

Early October 1989

- Hadronic event selection based on calorimeters
- Immense effort with many sleepless nights as we were at the same time validating the data, removing the noise and calibrating them and running the reconstruction

SELECTION OF HADRONIS / Z FOR  
CROSS-SECTION MEASUREMENT

S. Dugeay, F. Fidecaro, N. Minod, F. Palau  
M. Pepe, J. Steinberger

IN CLOSE CONTACT WITH E. BLUCHER, F. BIRD & LL. LARRID

■ OFFLINE TRIGGER BASED ON ECAL WIRES



6 GeV IN BARREL .OR. 3 GeV IN ENDCAPS IN  
COINCIDENCE


$$\epsilon_{q\bar{q}} = 99.8\%$$

$$\epsilon_{ZZ} = 74.1\%$$

$$\epsilon_{\gamma\gamma} = 2.3\%$$

■  $E_{\text{ECALW}} + E_{\text{HCAL}} > 20 \text{ GeV}$

# First ALEPH Z line shape



# CERN Theory Christmas play 1989



<https://videos.cern.ch/record/1337777>

# Foundation of the LEP ElectroWeak Working Group

- Originally, a group with members of the four LEP experiments, led by **Jack Steinberger**, investigated the combination of the Z line shape parameters [**Phys. Lett. B 276 (1992) 247**]
  - Jack insisted that the combination was a job for the experimentalists from the four collaborations who should discuss together, rather than for the theorists! [J.Lefrancois reminded me of Jack's role in this!]
- This led to the establishment of the **LEP ElectroWeak Working Group**, an **unprecedented**, collaborative effort across the experiments
- Mandate: to combine the measurements of the four LEP experiments on electroweak observables, e.g. cross sections, masses and various couplings, properly taking into account the common systematic uncertainties and producing the "best" LEP averages

<http://lepewwg.web.cern.ch/LEPEWWG/>

# Foundation of

- Originally, a group of physicists studying the **LEP EW parameters** [Physics Letters B 250 (1990) 155]
  - Jack insisted on the four collaboration theorists!
- This led to the **LEP EW Group**, an **unofficial** group of experimentalists
- Mandate: to coordinate the LEP experiments on electroweak couplings, provide a common set of parameters and uncertainties and



# Working Group


experiments, led by the Z line shape  
experimentalists from other than for the

**LEP EW Working Group**  
across the

LEP experiments assess and various systematic errors

<http://lepewwg.web.cern.ch/LEPEWWG/>

# Final combined result



$$N_{\nu} = 2.9840 \pm 0.0082$$



- Based on 17 million Z decays
- Less than 3 per mille uncertainty
- ~ half of it from theoretical uncertainty on low-angle Bhabha scattering cross-section

LEP EW WG:  
Phys. Rept. 427 (2006)

- First paper ever signed by over 2500 authors !



# An alternative method: $e^+e^- \rightarrow \nu\bar{\nu}\gamma$

- Detect events with a single photon and nothing else at energies above the Z mass (method originally advocated for  $\nu$  counting)
- Cross-section approximately proportional to  $N_\nu$  (contribution from  $t$ -channel  $W$ -exchange is small)
- Around 2500 single-photon events collected by the four LEP experiments, giving  $N_\nu = 3.00 \pm 0.08$
- By also including data at  $130 < \sqrt{s}$  [GeV]  $< 209$  for new physics searches, the LEP experiments collectively detected  $\sim 6200$  single-photon events, giving  $N_\nu = 2.92 \pm 0.005$



# $e^+e^- \rightarrow \nu\bar{\nu}\gamma$ (The return)

- Method advocated in FCC-ee studies for a high precision measurement of invisible partial width (sensitive to invisible particles, e.g. a neutralino, or to the mixing of heavy right-handed neutrinos with existing ones)
- Measure  $\Gamma_Z^{\text{inv}}/\Gamma_Z^{\text{lept}}$  above the Z to eliminate many systematic uncertainties (e.g. luminosity, photon detection efficiency)
- Gain of a factor  $\sim 10$  in precision on  $N_\nu$  seems possible



[<https://arxiv.org/pdf/1308.6176.pdf>]

# Conclusions

- The measurement of the number of neutrino generations stands out as one of the legacies of the LEP physics programme. It ruled out for the first time the existence of a fourth generation, posing stringent limits on theoretical models relevant in astrophysics and cosmology
- The overall determination of the Standard Model parameters by the LEP experiments, with precision exceeding the initial expectations, and the proof of its unexpected consistency, marked a turning point in our field
- The story of success of the Standard Model continues with the results from the LHC, demonstrating its validity up to the multi-TeV range and possibly even beyond
- Still many questions remain unanswered: Why are there just three families of particles? What determines the masses of their members? These questions still lie at the centre of particle physics today