

Recent Results from the IceCube Neutrino Observatory



Sanjib Kumar Agarwalla
sanjib@iopb.res.in



Institute of Physics, Bhubaneswar, India
(For the IceCube Collaboration)



The Nobel Prize in Physics 2002



Raymod Davis Jr.

Detected Solar Neutrinos




Masatoshi Koshihara

Detected Supernova Neutrinos


Detection of Cosmic Neutrinos → A New Window on the Universe

Era of Low-Energy Neutrino Astronomy began!

Detection of High-Energy Cosmic Neutrinos



Highest Energy Radiation from Universe: Neutrinos & Cosmic Rays



Universe beyond our Galaxy is opaque to gamma rays
Neutrinos open a new window to observe the high-energy and distant Universe

For a recent review see: Arguelles, Halzen, Kurahashi, arXiv: 2405.17623 [hep-ex]

Multi-messenger Connection

Accelerated cosmic ray beam (p) interact with gas (p) or radiation (γ).


$$p + p \rightarrow n_\pi [\pi^0 + \pi^+ + \pi^-] + X$$

$$p + \gamma \rightarrow \Delta^+ \rightarrow \begin{cases} p + \pi^0 & (2/3) \\ n + \pi^+ & (1/3) \end{cases}$$

$$\begin{cases} \pi^+ \rightarrow \nu_\mu + \mu^+ \rightarrow \nu_\mu + (e^+ + \nu_e + \bar{\nu}_\mu) \\ \pi^- \rightarrow \bar{\nu}_\mu + \mu^- \rightarrow \bar{\nu}_\mu + (e^- + \bar{\nu}_e + \nu_\mu) \\ \pi^0 \rightarrow \gamma + \gamma \end{cases}$$

Neutrinos are “smoking gun” of cosmic ray accelerators

Correlated neutrinos~gamma rays~cosmic rays emission rate at source



$$\frac{1}{3} \sum_{\nu_\alpha} E_\nu^2 Q_{\nu_\alpha} \simeq \frac{K_\pi}{4} E_\gamma^2 Q_\gamma \simeq \frac{1}{4} f_\pi \frac{K_\pi}{1 + K_\pi} E_p^2 Q_p$$


$$E_\nu \simeq \frac{1}{2} E_\gamma \simeq \frac{1}{20} E_p$$

One could use the CR flux to set a bound on the neutrino flux – Waxman-Bahcall bound for optically thin sources


$$E_\nu^2 I_\nu(E_\nu) \sim 5 \times 10^{-8} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1} \quad \text{Gigaton detector!}$$

Neutrinos open a new window to observe the high-energy and distant Universe


IceCube at South Pole



IceCube Neutrino Telescope





- 1 km³ ν detector deep under ice at South Pole
- 3 components: IceTop, IceCube, and DeepCore
- 5160 DOMs across 86 strings
- Optimized for TeV-PeV



- Neutrino interacts with ice and produces charged lepton
- Lepton direction closely aligned with neutrino
- Charged leptons emit Cherenkov radiation, when they travel faster than light in a medium
- Radiation detected by DOM (Digital Optical Modules)

Event Signatures in IceCube

IceCube detects neutrinos by measuring the Cherenkov light emitted by charged secondary particles from neutrinos interacting with the ice and the Antarctic bedrock




Cascade event

ν_e - CC (charged current) and
all flavor NC (neutral current)
interactions



Good energy resolution $\sim 15\%$
Poorer angular resolution $\sim 10^\circ$ ($\gtrsim 100$ TeV)




Track

ν_μ - CC and atmospheric muons



Poorer energy resolution \sim factor of 2
Excellent angular resolution $\sim 0.5^\circ$




Double-Bang

ν_τ - CC interactions



Distinctive double-cascade signatures
in the full detector

IceCube Science – A Multidisciplinary Instrument



RESEARCH

RESEARCH ARTICLES


Science June 30, 2023

NEUTRINO ASTROPHYSICS


Observation of high-energy neutrinos from the Galactic plane

IceCube Collaboration*†

DOI: [10.1126/science.adc9818](https://doi.org/10.1126/science.adc9818)




The Multiwavelength Milky Way



$\pm 10^\circ$ in latitude, from radio to gamma ray


S. K. Agarwalla, PASCOS 2024, ICISE, Quy Nhon, Vietnam, 8th July 2024

The Multiwavelength Multimessenger Milky Way



First Observation of the Diffuse Galactic Plane in Neutrinos!

Important Breakthrough in Selection of Astrophysical Neutrinos



Event rates in the past

Atmospheric muons: ~ 2700 per second
Atmospheric neutrinos: ~ 1 per hour
Astrophysical neutrinos: ~ 1 per day

Event selection at present

Employs series of convolutional neural networks (CNNs) & boosted decision trees

Improved reconstruction resolution over entire energy range

Huge improvement in cascade events

30 times as many events as precursor analysis


Background reduced by almost 8 orders of magnitude ($\sim 2700/\text{s}$ to $\sim 17/\text{day}$)

Analysis sensitivity improved by a factor of 3


Equivalent to savings of 75 years of detector lifetime > \$500 million

Improved Astronomy due to Cascade Events

Skymap from 2017 paper




Skymap from 2019 paper



2 years of data
Galactic Plane p-value: 65%

7 years of data
Galactic Plane p-value: 2.1% (2σ)



10 years of data
Galactic Plane p-value: 0.0004% (4.5σ)

Courtesy Naoko Kurahashi Neilson (ICRC 2023)

Take Home Message

+ Strong evidence for neutrino emission from the Galactic plane

- Background-only hypothesis rejected at 4.5σ
- Emission from Galactic plane may explain up to $\sim 10\%$ of astrophysical flux observed by IceCube
- Independent hints in IceCube track channels ($\sim 2.7\sigma$) and in ANTARES ($\sim 2\sigma$)

+ Observation enabled by new tools based on Deep Learning

- 30 times as many events than precursor selection
- Improved reconstruction resolution by up to 50%
- Analysis sensitivity improved by a factor of 3

+ This result leads to many new questions:

- Pure diffuse emission, or point sources in there too? Emission from the Sgr A*? What is the energy spectrum? Comparison with multiwavelength emissions...Origin of CRs? Galactic structure?.....

+ Hope:

- Ongoing studies, future upgrades, and comb. w/ other ν detectors may help to address these issues


We have arrived in the era of high-energy neutrino astronomy!

The First Steady Neutrino Source: NGC 1068

AGN promising candidate for neutrino source since 1970s

IceCube identified neutrino emission from NGC 1068, a Type II Seyfert galaxy, at 4.2σ in 2022 \Rightarrow very close, $z = 0.004$ (14 Mpc)


Soft best-fit spectrum power-law with spectral index $\gamma = 3.2 \pm 0.2$



IceCube, PRL 124, 051103 (2020)


The First Transient Source: TXS 0506+056

Multimessenger observations of a flaring blazar



IceCube-170922A 290 TeV neutrino
coincident with the TXS0506+056 blazar


Science 36, eaat1378 (2018)



Following an alert sent by IceCube on 23.09.2017, Fermi-LAT & MAGIC detected γ -flaring activity and very-high-energy γ rays, respectively, in the direction of blazar TXS 0506-056

IceCube investigated models associating neutrino & γ -ray production & find that correlation of the neutrino with the flare of TXS 0506+056 is statistically significant at 3σ

Astrophysical Neutrinos: Extragalactic Diffuse Flux



So far IceCube has identified three neutrino sources, but there are many open questions:

1. TXS 0506+056 blazar



2. NGC 1068 AGN



3. Galactic Plane








The majority of the astrophysical flux that IceCube sees comes from a diffuse component that remains to be fully understood, as it differs depending on the chosen event sample, sky coverage, energy range

Break from a single power-law?

Glashow Resonance: A prediction, 60 years later observed


- Resonant interaction $\bar{\nu}_e + e^- \rightarrow W^- \rightarrow X$
- Disentangle antineutrinos from the total neutrino flux.



Astrophysical Neutrinos: Flavor Ratios




pion production




(1:2:0)

neutron decay





(1:0:0)

muon dumped



(0:1:0)



Eur. Phys. J. C 82, 1031 (2022)

Courtesy J. A. Aguilar

An important tool to search for the BSM physics in IceCube

See talk by Kareem Farrag in this conference

Seven Astrophysical Tau Neutrino Candidates

PHYSICAL REVIEW LETTERS 132, 151001 (2024)

Editors' Suggestion

Featured in Physics

Observation of Seven Astrophysical Tau Neutrino Candidates with IceCube



R. Abbasi,¹⁷ M. Ackermann,⁶² J. Adams,¹⁸ S. K. Agarwalla,^{40,*} J. A. Aguilar,¹² M. Ahlers,²² J. M. Alameddine,²³ N. M. Amin,⁴⁴ K. Andeen,⁴² G. Anton,²⁶ C. Argüelles,¹⁴ Y. Ashida,⁵³ S. Athanasiadou,⁶² S. N. Axani,⁴⁴ X. Bai,⁵⁰ V. A. Balagopal,⁴⁰ M. Baricevic,⁴⁰ S. W. Barwick,³⁰ V. Basu,⁴⁰ R. Bay,⁸ J. J. Beatty,^{20,21} J. Becker Tjus,^{11,†} J. Beise,⁶⁰ C. Bellenghi,²⁷ C. Benning,¹ S. BenZvi,⁵² D. Berley,¹⁹ E. Bernardini,⁴⁸ D. Z. Besson,³⁶ E. Blaufuss,¹⁹ S. Blot,⁶²

(IceCube Collaboration)[§]

additional authors not shown

We report on a measurement of astrophysical tau neutrinos with 9.7 yr of IceCube data. Using convolutional neural networks trained on images derived from simulated events, seven candidate events were found with visible energies ranging from roughly 20 TeV to 1 PeV and a median expected parent ν_τ energy of about 200 TeV. Considering backgrounds from astrophysical and atmospheric neutrinos, and muons from π^\pm/K^\pm decays in atmospheric air showers, we obtain a total estimated background of about 0.5 events, dominated by non- ν_τ astrophysical neutrinos. Thus, we rule out the absence of astrophysical ν_τ at the 5 σ level. The measured astrophysical ν_τ flux is consistent with expectations based on previously published IceCube astrophysical neutrino flux measurements and neutrino oscillations.

DOI: 10.1103/PhysRevLett.132.151001



Detection of astrophysical ν_τ require shower-like bright events (double cascades)


Study images of 3 brightest neighbouring strings

Train 3 CNNs to distinguish astrophysical ν_τ from possible backgrounds

7 astrophysical ν_τ candidates identified with CNN using 9.7 years of IceCube data – some events show clear double-pulse waveforms

We rule out the absence of astrophysical ν_τ at 5.1 σ


DeepCore Detector



- 8 dedicated strings with denser spacing
- 40–70 m horizontally & 7 m vertically
- Optimized for GeV scale neutrinos
- Uses IceCube as VETO
- Fiducial volume ~ 10 Mton

The design and performance of IceCube DeepCore (2012): [Astroparticle Physics, 35\(10\), 615-624 \(2012\)](#)

Oscillograms for Muon Neutrino Survival Channel



Parametric resonance region

$\cos\theta_\nu < -0.8$
 $3 \text{ GeV} < E_\nu < 6 \text{ GeV}$
reducing threshold helps

Useful to probe
Earth's matter effect


MSW resonance region

$-0.8 < \cos\theta_\nu < -0.5$
 $6 \text{ GeV} < E_\nu < 10 \text{ GeV}$

MSW or parametric resonances have
not been observed yet inside Earth!


Simulated Neutrino Events

- Convolutional Neural Network (CNN) based reconstruction
- Monte Carlo (MC) simulation with 9.3 years of exposure (2012 - 2021)
- Huge statistics (~192k events)
- Neutrinos comprise 99.5% of sample
- High statistics in ν_μ CC channel
- Filters are applied to eliminate primary backgrounds: noise and atm. muon contamination (~0.5%)





Selection	Expected MC Events (9.3 yr)	% of Sample
$\nu_e + \bar{\nu}_e$ CC	48616	25.2
$\nu_\mu + \bar{\nu}_\mu$ CC	110656	57.5
$\nu_\tau + \bar{\nu}_\tau$ CC	10938	5.7
$\nu_{\text{all}} + \bar{\nu}_{\text{all}}$ NC	21412	11.1
μ_{atm}	973	0.5
All MC	192597	—

Latest Oscillation Results from DeepCore



[arXiv:2405.02163](https://arxiv.org/abs/2405.02163)


- These new result are compatible & complementary with the existing measurements
 - High-precision measurement on Δm^2_{32}
- Comparatively high-energy sample (5 – 100 GeV) and different systematic uncertainties
 - Strong validation of the standard 3-flavor oscillation



[ICRC2023 arXiv:2307.15855](https://arxiv.org/abs/2307.15855)


Sensitivity towards Neutrino Mass Ordering and Earth's Matter Effect

DeepCore Neutrino Mass Ordering (9.28 years)




See talks given in NuFact 2023 and NuTel 2023

12-layered PREM (True) vs. Vacuum (Test)





See talk given in Brookhaven Forum 2023

Validating Broad Features of the PREM Profile of Earth





See talks given in Brookhaven Forum 2023





A New Extension of DeepCore: IceCube Upgrade

- 2 Mton of dense instrumentation for low-energy measurements
- 7 new strings in the center region of detector: energy threshold ~ 1 GeV
- Higher event rate: 4 x DeepCore
- To be deployed in the Antarctic summer of 2025/26



30 GeV Neutrino

Spacing between new modules:
20 m horizontally & 3 m vertically





2 new types of optical modules w/ multi-PMT configurations

[ICRC2019 arXiv:1908.09441](#)

[ICRC2023 arXiv:2307.15295](#)

Sensitivity of IceCube Upgrade: Atmospheric Oscillation Parameters





[ICRC2023 arXiv:2307.15295](#)

- 90% confidence level after 3 years with the new strings assuming NuFit 5.2 best-fit values
- With the new strings, IceCube's sensitivity to Δm_{31}^2 and θ_{23} increases by about 20 to 30%
- 4 times enhancement in the sensitivity to neutrino mass ordering




IceCube-Gen2: A Massive Telescope

From IceCube-Gen2 Collaboration (2023): IceCube-Gen2 Technical Design: The IceCube-Gen2 Neutrino Observatory

<https://icecube-gen2.wisc.edu/science/publications/TDR>



IceCube-Gen2 is the proposed extension of IceCube detector: increase the instrumented volume from 1 km³ to ~ 8 km³, 9600 additional modules, extend to higher energy (10¹⁸ eV) with radio and surface array, and new detector design



Proposed detector designs based on mDOM and D-Egg from the Upgrade

Concluding Remarks

- Over the past decade, the IceCube Neutrino Observatory has opened up a new window onto the extreme and hidden universe. It has detected high-energy neutrinos of astrophysical origin and successfully identified the first sources.
- The DeepCore array in the central region of IceCube has enabled the detection and reconstruction of atmospheric neutrinos with energies as low as a few GeV, providing high-precision measurements of oscillation parameters and first glimpse of Earth matter effects.
- IceCube Upgrade - a new low-energy extension of DeepCore with novel sensors will address several open questions in three-flavour neutrino oscillation paradigm and significantly improve the detector calibration, resulting in better particle identification, angular, and energy resolution.
- The proposed IceCube-Gen2 will substantially increase the detection rate of astrophysical neutrinos with highest energies to better understand their possible sources and source populations.

Thank you!